

Dark soliton oscillations in Bose-Einstein condensates with multi-body interactions

Anatoly M Kamchatnov¹ and Mario Salerno²

¹ Institute of Spectroscopy, Russian Academy of Sciences, Troitsk, Moscow Region, 142190, Russia

² Dipartimento di Fisica “E. R. Caianiello”, Consorzio Nazionale Interuniversitario per le Scienze Fisiche della Materia (CNISM), Università di Salerno, via S. Allende I-84081, Baronissi (SA), Italy

E-mail: kamch@isan.troitsk.ru, salerno@sa.infn.it

Abstract. We consider the dynamics of dark matter solitons moving through non-uniform cigar-shaped Bose-Einstein condensates described by the mean field Gross-Pitaevskii equation with generalized nonlinearities, in the case when the condition for the modulation stability of the Bose-Einstein condensate is fulfilled. The analytical expression for the frequency of the oscillations of a deep dark soliton is derived for nonlinearities which are arbitrary functions of the density, while specific results are discussed for the physically relevant case of a cubic-quintic nonlinearity modeling two- and three-body interactions, respectively. In contrast to the cubic Gross-Pitaevskii equation for which the frequencies of the oscillations are known to be independent of background density and interaction strengths, we find that in the presence of a cubic-quintic nonlinearity an explicit dependence of the oscillations frequency on the above quantities appears. This dependence gives rise to the possibility of measuring these quantities directly from the dark soliton dynamics, or to manage the oscillation via the changes of the scattering lengths by means of Feshbach resonance. A comparison between analytical results and direct numerical simulations of the cubic-quintic Gross-Pitaevskii equation shows good agreement which confirms the validity of our approach.

PACS numbers: 03.75.Kk

Submitted to: *J. Phys. B: At. Mol. Phys.*

1. Introduction

Since the first experimental realization of Bose-Einstein condensates (BECs) of diluted atomic gases, the dynamics of matter waves in different trapping potentials has attracted the attention of many researchers and presently is a field of growing interest [1, 2]. The dispersive properties of BEC induced by the quantum properties of a slow atomic motion, on one side, and the intrinsic nonlinearity of the system induced by the interatomic interactions, on the other side, make BECs ideal systems for exploring competing effects of dispersion and nonlinearity. This results in a variety of spectacular phenomena ranging from the existences of matter wave vortices [3] and solitons [4, 5, 6] to dispersive shocks [7, 8].

In the standard mean field approximation [1] the BEC dynamics is governed by the Gross-Pitaevskii (GP) equation

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \Delta \psi + V(\mathbf{r})\psi + f(|\psi|^2)\psi \quad (1)$$

for the condensate “wave function” ψ , where $V(\mathbf{r})$ is the potential of the external forces trapping the condensate. In the standard GP approximation

$$f(|\psi|^2) = g|\psi|^2, \quad (2)$$

where $g = 4\pi\hbar^2 a_s/m$ is the nonlinear interaction parameter expressed in terms of the s -wave scattering length a_s of atoms with mass m . In this approximation, only two-body collisions are taken into account. The scattering length a_s , however, can be varied and even made equal to zero by means of Feshbach resonances. Then contribution of multi-particle (three-body) collisions becomes comparable with the two-body contribution and a more complicated form of the nonlinear term, such as

$$f(|\psi|^2) = g|\psi|^2 + h|\psi|^4 \quad (3)$$

arises in the GP equation. Although the three-dimensional GP equation (1) is too complicated for analytical studies, one can often consider experimental low dimensional settings, e.g. cigar-shaped traps with a strong radial confinement, for which the radial motion of the BEC is “frozen” and Eq. (1) reduces to an effective 1D cubic-quintic generalized nonlinear Schrödinger (NLS) equation [9],

$$i\psi_t = -\frac{1}{2}\psi_{xx} + V(x)\psi + f(|\psi|^2)\psi, \quad (4)$$

(here standard non-dimensional variables have been introduced, holding the previous notation for convenience). In this reduction, the function $f(\rho)$, $\rho = |\psi|^2$, preserves its form (3) up to a renormalization of the constants g and h . Also note that the conservation of the norm of the condensate wave function (normalized number of atoms) is guaranteed for arbitrary functions $f(\rho)$. More complicated form of the function $f(\rho)$ can arise if we take into account the radial motion of BEC (see, e.g., [10, 11]). Thus, we arrive at the problem of 1D dynamics of BEC with generalized form of the nonlinearity $f(\rho)$ and confined in the axial direction x by the trap $V(x)$.

As is well known, in case of $f(\rho)$ given by Eq. (2) with $g > 0$ (repulsive two body interaction) and under the assumption of negligible changes of the potential $V(x)$ in the region of x under consideration, the NLS equation (4), (2), has a dark soliton solution corresponding to the motion of a hole in the density distribution function, $\rho(x - vt)$, along a uniform distribution $\rho = \rho_0$ with constant velocity v which depends on the depth of the soliton [12]. If the potential $V(x)$ is a slowly varying function for distances of the order of the soliton's width, then the coordinate of the soliton is well defined and we can speak about value of the potential $V(x)$ at the “point” x of the soliton position. In this case we can formulate the problem of motion of a dark soliton “in the potential” $V(x)$ in terms of its collective coordinate. This problem has been addressed in several papers (see, e.g., [13, 14, 15, 16, 17, 18, 19]) where it was shown that the dynamics of dark solitons is quite nontrivial. In particular, if a BEC described by the standard GP equation (2) is confined in a harmonic axial trap,

$$V(x) = \frac{1}{2}\omega_0 x^2, \quad (5)$$

then the dark soliton oscillates inside the condensate with the frequency [13]

$$\omega = \frac{\omega_0}{\sqrt{2}}. \quad (6)$$

This result is quite surprising because apparently it contradicts to a simple consequence of the Ehrenfest theorem which says that the center of mass of the whole BEC always oscillates with the trap frequency ω_0 . The remarkable theoretical prediction (6) was confirmed in the experiment [20]. Equation (6) then implies that the motion of the dark soliton must be accompanied by a deformation of the density distribution which re-tunes the oscillation of the whole condensate to the trap frequency. In this situation the motion of the dark soliton becomes “decoupled” from the motion of the center of mass and the Ehrenfest theorem is not violated.

However, the result (6) is not universal. If the nonlinearity function is given by

$$f(\rho) = h\rho^2 \quad (7)$$

with $h > 0$, then the frequency ω of oscillations of a shallow dark soliton coincides with the trap frequency [14],

$$\omega = \omega_0 \quad (8)$$

and for deep solitons this frequency is given by a more complicated expression (see [16])

$$\omega = \left\{ 1 + \frac{\sqrt{3}}{2 \ln \frac{1+\sqrt{3}}{\sqrt{2}}} \right\}^{-1/2} \omega_0 \approx 0.6572\omega_0. \quad (9)$$

This observation poses the problem of the dark soliton motion in the case of a more general form of the nonlinearity.

The aim of the present paper is to address this problem by considering a non-uniform cigar-shaped BEC with nonlinear interactions modeled by an arbitrary function $f(\rho)$ of the density. In particular we consider the physically relevant example of the

cubic-quintic nonlinearity function in (3), modeling the mean field effect of the two-body and three-body collisions in the GP equation. In this case we find that the relative contribution of the two nonlinear terms to the soliton oscillation depends on the background density ρ_0 : if a deep soliton oscillates in vicinity of the maximum ρ_0 of the density distribution, then for $|h|\rho_0 \ll g$ its frequency must be equal to (6), and for $|h|\rho_0 \gg g$ to (9). The transition from one limiting case to another is of considerable theoretical interest. We derive an analytical expression for the oscillation frequency of a deep dark solitons which reveals the explicit dependence of the oscillations frequency both on the background density and on the strength of the nonlinearity parameters. Our formulae reproduce the results considered in [15, 16] for the particular cases (2) and (7) of the nonlinearity for which the above dependence disappears.

We also consider the case of a quintic attractive nonlinearity ($h < 0$) with an overall repulsive interaction. In this case for the density $\rho > g/(2|h|)$ we get modulation instability of BEC, i.e. the sound velocity

$$c_s = \sqrt{\rho_0 f'(\rho_0)} \quad (10)$$

becomes imaginary so that the notion of “dark solitons” loses its sense. We show that in this case the oscillation frequency ω as a function of ρ_0 becomes singular at $\rho_{0c} = g/(2|h|)$. The form of this singularity may provide further advances in the understanding of the dynamics of dark solitons in BEC.

Our theory will be based on the elegant approach of Refs. [15, 16] where the dark soliton was considered as a “Landau quasi-particle” whose motion in the stationary potential $V(x)$ is governed by the quasi-particle’s “energy conservation” law in close analogy with the Newtonian particle motion. The generalization of this approach to arbitrary $f(\rho)$ allows us to derive the equation of motion of the dark soliton for the general case and to show the dependence of the soliton effective mass on the system parameters. A comparison between analytical results and direct numerical simulations of the cubic-quintic GP equation shows a very good agreement which further confirms the validity of our analysis.

We finally remark that the dependence of the dark soliton oscillation frequency on the system parameters can be very useful for experimental studies of relative contribution of multi-particle collisions into BEC dynamics by varying the value of g by means of Feshbach resonance. Moreover, it gives the possibility to measure background densities or scattering lengths directly from the dark soliton dynamics. The possibility to manage soliton oscillations via temporal changes of the scattering lengths is also an attractive feature emerging from our study which could be of interest for practical applications.

The paper is organized as follows. In Section II we discuss the general theory of the dynamics of a dark soliton of the NLS equation with an arbitrary nonlinearity. The results will be used to study the cases of both deep and shallow soliton dynamics. In Section III we consider oscillations of dark solitons for the specific case of a cubic-quintic nonlinearity. We consider the cases of a repulsive cubic nonlinearity and either

repulsive or attractive quintic nonlinearity, the last being done for parameters satisfying the condition of modulation stability of BEC. In Section IV analytical results will be compared with direct numerical simulations of the cubic-quintic GP equation for the case of repulsive and attractive cubic and quintic nonlinearity, respectively. In the last Section we draw our conclusions and summarize the main results of the paper.

2. Motion of a dark soliton in a trap: general theory

First, we have to find the soliton solution of the generalized NLS equation (4) with zero potential $V(x) \equiv 0$ but under the boundary condition that the condensate density $\rho(x)$ tends to the limiting background density ρ_0 as $|x| \rightarrow \infty$ and the condensate's velocity vanishes at infinity. To this end, it is convenient to look for the solution in the form

$$\psi = \sqrt{\rho(\xi)} \exp(i\varphi(\xi) - i\mu t) \quad (11)$$

where

$$\mu = f(\rho_0) \quad (12)$$

is the chemical potential of the condensate and its density ρ and the local flow velocity $u = \varphi_x$ depend only on the variable $\xi = x - vt$ with v being the soliton velocity. The formulated above boundary conditions take the form

$$\rho \rightarrow \rho_0, \quad \varphi_\xi \rightarrow 0, \quad \text{as } |\xi| \rightarrow \infty. \quad (13)$$

Substitution of (11) into (4) with $V(x) \equiv 0$ and separation of real and imaginary parts yields the system of two equations for $\rho(\xi)$ and $u(\xi) = \varphi_\xi$. One of these equations can be integrated at once to give

$$u(\xi) = \varphi_\xi = v \left(1 - \frac{\rho_0}{\rho} \right) \quad (14)$$

where the integration constant is chosen according to the conditions (13). Then the second equation can be transformed to

$$\frac{1}{8}\rho_\xi^2 - \frac{1}{4}\rho\rho_{\xi\xi} + \rho^2 f(\rho) - \mu\rho^2 + \frac{1}{2}v^2(\rho_0^2 - \rho^2) = 0 \quad (15)$$

and this equation can also be integrated once. As a result we obtain the equation

$$\rho_\xi^2 = Q(\rho) \quad (16)$$

where

$$Q(\rho) = 8\rho \int_\rho^{\rho_0} [\mu - f(\rho')] d\rho' - 4v^2(\rho_0 - \rho)^2 \quad (17)$$

and again the integration constant is chosen according to the condition (13). Thus, we have reduced the problem of finding the density profile to the inversion of the integral

$$\xi = \int_{\rho_m}^\rho \frac{d\rho}{\sqrt{Q(\rho)}} \quad (18)$$

where ρ_m is the minimal density at the center $\xi = 0$ of the soliton. The function $Q(\rho)$ has a double zero at $\rho = \rho_0$ and the condition $dQ(\rho)/d\rho|_{\rho=\rho_0} = 0$ reproduces the

relation (12). At $\rho = \rho_m$ the function $Q(\rho)$ has a single zero, and this condition yields the relationship between ρ_m and the soliton velocity v ,

$$v^2 = \frac{Q_0(\rho_m)}{4(\rho_0 - \rho_m)^2}, \quad (19)$$

where

$$Q_0(\rho) = 8\rho \int_{\rho}^{\rho_0} [f(\rho_0) - f(\rho')] d\rho' \quad (20)$$

is the function (17) with zero value of the velocity v and we have used here Eq. (12) for μ .

Now we have to obtain the expression for the energy of the dark soliton. To this end, we notice that the function $\Psi = \sqrt{\rho} \exp(i\varphi) = \psi \cdot \exp(i\mu t)$ satisfies the equation

$$i\Psi_t + \frac{1}{2}\Psi_{xx} + (\mu - f(|\Psi|^2))\Psi = 0 \quad (21)$$

which can be written in a Hamiltonian form

$$i\Psi_t = \frac{\delta H}{\delta \Psi^*} \quad (22)$$

with

$$H = \int_{-\infty}^{\infty} \left[\frac{1}{2} |\Psi_x|^2 + \int_0^{|\Psi|^2} f(\rho) d\rho - \mu |\Psi|^2 \right] dx. \quad (23)$$

To get the contribution of the dark soliton into the energy of BEC, we subtract the background energy $H_0 = \int_{-\infty}^{\infty} \left[\int_0^{\rho_0} f(\rho) d\rho - \mu |\Psi|^2 \right] dx$ and express the resulting formula in terms of ρ and φ_x ,

$$E = H - H_0 = \int_{-\infty}^{\infty} \left[\frac{1}{2} \left(\varphi_x^2 + \frac{\rho_x^2}{4\rho^2} \right) \rho + \int_{\rho}^{\rho_0} [f(\rho_0) - f(\rho')] d\rho' \right] dx. \quad (24)$$

Substitution of expressions (14) and (16) for the dark soliton solution casts this formula into

$$E = 2 \int_{-\infty}^{\infty} \int_{\rho}^{\rho_0} [f(\rho_0) - f(\rho')] d\rho' dx. \quad (25)$$

At last, integration with respect to x can be transformed to integration with respect to ρ from ρ_m to ρ_0 with account of (18),

$$E = 4 \int_{\rho_m}^{\rho_0} \frac{d\rho}{\sqrt{Q(\rho)}} \int_{\rho}^{\rho_0} [f(\rho_0) - f(\rho')] d\rho' = \frac{1}{2} \int_{\rho_m}^{\rho_0} \frac{Q_0(\rho)}{\rho \sqrt{Q(\rho)}} d\rho. \quad (26)$$

Here ρ_m can be considered according to (19) as a function of ρ_0 and v^2 , hence the soliton energy is also a function of these two variables,

$$E = E(\rho_0, v^2). \quad (27)$$

Now we turn to the problem of motion of the dark soliton along a nonuniform condensate confined in a trap with the axial potential $V(x)$. Then in the Thomas-Fermi (TF) approximation the density $\rho(x)$ is a function of x determined implicitly by the equation

$$\mu - V(x) = f(\rho(x)). \quad (28)$$

When this TF distribution of the density $\rho = \rho(x)$ is substituted into (27) instead of ρ_0 , we obtain the energy of the soliton moving with velocity v and located at this moment at the point x . As was shown in Ref. [15], the velocity v changes during the motion of the dark soliton in such a way that the soliton energy (27) is conserved. If we denote the soliton coordinate as $x = X(t)$, then $v = \dot{X}(t)$ and Eq. (27) converts into

$$E(\rho(X), \dot{X}^2) = E_0, \quad (29)$$

where E_0 is the initial energy of the soliton. Since the function $E(\rho, v^2)$ is known, Eq. (29) is the differential equation for finding the soliton coordinate X as a function of time t . In general, this first order ordinary differential equation should be solved numerically. Here we shall consider the most interesting limiting cases admitting complete analytical study.

2.1. Small amplitude oscillations of a deep soliton

Let us consider oscillations of a deep soliton,

$$\rho_m \ll \rho_0, \quad (30)$$

where ρ_0 is the maximal density of the condensate at its “top” when it is confined in a trap with the potential $V(x)$ (see (28)). In this case the amplitude of oscillations is much less than the TF radius of the condensate and the velocity of its motion is much less than the local sound velocity

$$v \ll c_s, \quad c_s = \sqrt{\rho_0 f'(\rho_0)}. \quad (31)$$

Therefore, if we expand the energy into series with respect to small amplitude and velocity, then the first nontrivial terms in (29) will give the expression

$$\frac{1}{2} m_* \dot{X}^2 + V(X) = \text{const}, \quad (32)$$

where $V(X)$ changes little for the small amplitude vibrations. Hence, we get the “energy conservation law” for a Newtonian particle moving in the potential $V(X)$, where m_* is the effective mass of the dark soliton.

For small v we obtain from (19)

$$\rho_m \cong \frac{\rho_0^2 v^2}{2a} \quad (33)$$

where

$$a = \int_0^{\rho_0} [f(\rho_0) - f(\rho)] d\rho. \quad (34)$$

To estimate the first term of the series expansion of E with respect to v^2 , we split the integral (26) into two terms (see [16]),

$$E = E_1 + E_2, \quad E_1 = \frac{1}{2} \int_{\rho_m}^{\rho_1} \frac{Q_0(\rho)}{\rho \sqrt{Q(\rho)}} d\rho, \quad E_2 = \frac{1}{2} \int_{\rho_1}^{\rho_0} \frac{Q_0(\rho)}{\rho \sqrt{Q(\rho)}} d\rho, \quad (35)$$

where we have introduced an intermediate integration limit ρ_1 so that

$$\rho_m \ll \rho_1 \ll \rho_0. \quad (36)$$

Dependence of E_1 on v^2 results mainly from the lower integration limit (33) and in the main approximation we get

$$E_1 \cong \frac{1}{2} \int_{\rho_m}^{\rho_1} \sqrt{Q(\rho)} \frac{d\rho}{\rho}. \quad (37)$$

Now, both limits of integration are small and ρ_m is a simple zero of the function $Q(\rho)$, hence we can approximate it as

$$Q(\rho) \cong \left. \frac{dQ(\rho)}{d\rho} \right|_0 (\rho - \rho_m) = 8a(\rho - \rho_m), \quad (38)$$

where a is defined in (34). Then substitution of (38) into (37) and elementary integration yield

$$E_1 \cong \sqrt{8a(\rho_1 - \rho_m)}. \quad (39)$$

We need the coefficient of v^2 in the expansion of E in powers of v^2 and the contribution of E_1 into this coefficient is equal to

$$\frac{dE_1}{d\rho_m} \frac{d\rho_m}{dv^2} = -\frac{\rho_0^2}{\sqrt{2a\rho_1}}. \quad (40)$$

As we see, it diverges in the limit $\rho_1 \rightarrow 0$. To cancel this divergent term, we have to identify a similar term in the derivative dE_2/dv^2 given by the expression

$$\left. \frac{dE_2}{dv^2} \right|_{v=0} = \int_{\rho_1}^{\rho_0} \frac{(\rho_0 - \rho)^2}{\rho \sqrt{Q_0(\rho)}} d\rho. \quad (41)$$

According to Eq. (40), the diverging term is proportional to $\rho_1^{-1/2}$. Correspondingly, we transform the integral in (41) in the following way by integration by parts,

$$\begin{aligned} \left. \frac{dE_2}{dv^2} \right|_{v=0} &= \int_{\rho_1}^{\rho_0} \frac{(\rho_0 - \rho)^2}{\rho^{3/2} \sqrt{Q_0(\rho)/\rho}} d\rho = \int_{\rho_1}^{\rho_0} \frac{(\rho_0 - \rho)^2}{\sqrt{Q_0(\rho)/\rho}} d(-2\rho^{-1/2}) \\ &= -\left. \frac{2(\rho_0 - \rho)^2}{\sqrt{Q_0(\rho)}} \right|_{\rho_1}^{\rho_0} + \int_{\rho_1}^{\rho_0} \frac{2}{\sqrt{\rho}} \frac{d}{d\rho} \left[\frac{\sqrt{\rho}(\rho_0 - \rho)^2}{\sqrt{Q_0(\rho)}} \right] d\rho. \end{aligned} \quad (42)$$

The first term vanishes at the upper limit $\rho = \rho_0$ and at the lower limit $\rho = \rho_1 \ll \rho_0$ we can approximate $Q_0(\rho)$ as $Q_0(\rho) \cong 8a\rho$ (see (38)). In the second term the integral is convergent and we can take here the lower integration limit equal to $\rho_1 = 0$. As a result we get

$$\left. \frac{dE_2}{dv^2} \right|_{v=0} \cong \frac{\rho_0^2}{\sqrt{2a\rho_1}} + 2 \int_0^{\rho_0} \frac{d\rho}{\sqrt{\rho}} \frac{d}{d\rho} \left[\frac{\sqrt{\rho}(\rho_0 - \rho)^2}{\sqrt{Q_0(\rho)}} \right]. \quad (43)$$

The first term cancels with the contribution (40) and we obtain the expression for the soliton mass

$$m_s = 2 \left. \frac{dE_2}{dv^2} \right|_{v=0} = 4 \int_0^{\rho_0} \frac{d\rho}{\sqrt{\rho}} \frac{d}{d\rho} \left[\frac{\sqrt{\rho}(\rho_0 - \rho)^2}{\sqrt{Q_0(\rho)}} \right] \quad (44)$$

for the motion of a dark soliton near the “top” of the condensate.

Introduction of the potential $V(x)$ leads to the shift of the chemical potential $\mu \rightarrow \mu - V(x)$. Taking into account the thermodynamic formula

$$N = -\frac{dE}{d\mu} \quad (45)$$

and the expression for the energy in the limit $v^2 \rightarrow 0$,

$$\begin{aligned} E = E_0 &= 4 \int_0^{\rho_0} \frac{d\rho}{\sqrt{Q_0(\rho)}} \int_\rho^{\rho_0} [\mu - f(\rho')] d\rho' \\ &= 4 \int_0^{\rho_0} \frac{d\rho}{\sqrt{8\rho}} \left\{ \int_\rho^{\rho_0} [\mu - f(\rho')] d\rho' \right\}^{1/2}, \end{aligned} \quad (46)$$

hence

$$N = -2 \int_0^{\rho_0} \frac{(\rho_0 - \rho) d\rho}{\sqrt{Q_0(\rho)}}, \quad (47)$$

we find the expansion of the soliton energy as

$$E = E_0 + \left. \frac{dE}{dv^2} \right|_{v^2=0} \cdot v^2 + \frac{dE}{d\mu} \cdot (-V(x)) = E_0 + \frac{1}{2} m_s v^2 + NV(x), \quad (48)$$

and division by N yields the energy conservation law for the soliton's motion

$$\frac{m_*}{2} v^2 + V(x) = \text{const} \quad (49)$$

where

$$m_* = \frac{m_s}{N} \quad (50)$$

is the effective mass of the soliton's motion. Hence, in case of a harmonic trap (5) we get the formula

$$\omega = \frac{\omega_0}{\sqrt{m_*}} \quad (51)$$

for the frequency of oscillations of deep solitons.

For a usual GP equation we have

$$f(\rho) = \rho, \quad (52)$$

hence

$$Q_0(\rho) = 4\rho(\rho_0 - \rho)^2, \quad m_s = -4\sqrt{\rho_0}, \quad N = -2\sqrt{\rho_0} \quad (53)$$

and

$$m_* = 2 \quad (54)$$

so that (51) reproduces the well-known result.

2.2. Motion of a shallow soliton

In the shallow soliton limit

$$\rho_0 - \rho \ll \rho_0 \quad (55)$$

the series expansion of Eq. (19) gives

$$v^2 \cong \rho_m \left[f'(\rho_0) - \frac{1}{3} f''(\rho_0)(\rho_0 - \rho_m) \right] \quad (56)$$

or with account of the expression (10) for the sound velocity we obtain

$$\rho_m = \rho_0 - \frac{c_s^2 - v^2}{f'(\rho_0) + \frac{1}{3}\rho_0 f''(\rho_0)}. \quad (57)$$

In this limit the dark soliton can be considered in the Korteweg-de Vries (KdV) approximation (see, e.g. [21]) for which the NLS equation with $V(x) \equiv 0$ reduces to the equation

$$\rho'_t + c_s \rho'_x + \frac{3}{2c_s} \left[f'(\rho_0) + \frac{1}{3}\rho_0 f''(\rho_0) \right] \rho' \rho'_x - \frac{1}{8c_s} \rho'_{xxx} = 0 \quad (58)$$

for evolution of small deviations $\rho' = \rho_0 - \rho$ of the density from the background value ρ_0 . This equation has the soliton solution

$$\rho' = \rho_0 - \rho = \frac{c_s^2 - v^2}{f'(\rho_0) + \frac{1}{3}\rho_0 f''(\rho_0)} \cdot \frac{1}{\cosh^2[\kappa(x - vt)]} \quad (59)$$

where

$$\kappa = 2 \sqrt{\frac{c_s(c_s^2 - v^2)}{f'(\rho_0) + \frac{1}{3}\rho_0 f''(\rho_0)}} \quad (60)$$

is the inverse soliton's half-width and the minimal density ρ_m at the soliton's center $x = vt$ is given by the expression (57).

In the same approximation

$$Q(\rho) = 4 \left[f'(\rho_0) + \frac{1}{3}\rho_0 f''(\rho_0) \right] (\rho_0 - \rho)^2 (\rho - \rho_m) \quad (61)$$

and calculation of the soliton energy (26) gives

$$E = \frac{4}{3} \frac{f'(\rho_0)}{[f'(\rho_0) + \frac{1}{3}\rho_0 f''(\rho_0)]} \cdot (c_s^2 - v^2)^{3/2}. \quad (62)$$

If the width $1/\kappa$ of the soliton is much less than the condensate's size (the TF radius), then we can consider the motion of such a soliton in the trap by replacement of ρ_0 by the solution $\rho(x)$ of Eq. (28) and of v by its variable velocity dX/dt to get the equation

$$\left(\frac{dX}{dt} \right)^2 = \rho f'(\rho) - \left\{ \frac{3E}{4f'(\rho)} \left[f'(\rho) + \frac{1}{3} f''(\rho) \right]^2 \right\}^{2/3}, \quad (63)$$

where $\rho = \rho(X)$ is given by Eq. (28). For example, in case of $f = h\rho^2$, $h > 0$, we have $f'(\rho) = 2h\rho$ and $\mu - V(X) = h\rho^2$, hence (63) reduces to

$$\frac{1}{2} \left(\frac{dX}{dt} \right)^2 + V(X) + 2 \left(\frac{hE}{3} \right)^{2/3} \left(\frac{\mu - V(X)}{h} \right)^{1/3} = \mu. \quad (64)$$

As we see, for a harmonic trap (5) the frequency of oscillations depends on the energy E of the soliton. However, if E is small enough,

$$E \ll \frac{\mu}{\sqrt{h}}, \quad (65)$$

then the last term in the left-hand side can be neglected and we return to the usual energy conservation law for a Newtonian particle which for a harmonic trap gives Eq. (8) in agreement with [14, 16].

3. Oscillation of a dark soliton for the cubic-quintic nonlinearity

Now we shall consider in some detail the motion of a dark soliton described by the generalized NLS equation with $f(\rho)$ given by Eq. (3), i.e.,

$$f(\rho) = g\rho + h\rho^2. \quad (66)$$

Conditions for realization of motion of a shallow soliton are very restrictive and we shall confine ourselves to the most practical case of small oscillations of a deep soliton (Sec. 2A).

Substitution of (66) into (20) gives

$$Q_0(\rho) = 8\rho(\rho_0 - \rho)^2 \left[\frac{1}{2}g + \frac{1}{3}h(2\rho_0 + \rho) \right]. \quad (67)$$

Situations with $h > 0$ and $h < 0$ should be considered separately.

3.1. Repulsive quintic interaction ($h > 0$)

In this case elementary integration of Eq. (18) with $Q(\rho)$ given by (67) yields the profile of the density in the dark solitons solution

$$\rho(\xi) = \rho_0 - \frac{(\rho_0 - \rho_+)(\rho_0 - \rho_-)}{(\rho_+ - \rho_-) \cosh^2(\kappa\xi) + \rho_0 - \rho_+}, \quad \xi = x - vt, \quad (68)$$

where

$$\rho_{\pm} = \pm \sqrt{\left(\frac{3g}{4h} + \rho_0 \right)^2 + \frac{3v^2}{2h}} - \left(\frac{3g}{4h} + \rho_0 \right) \quad (69)$$

and

$$\kappa = \sqrt{2h(\rho_0 - \rho_+)(\rho_0 - \rho_-)}/3. \quad (70)$$

Substitution of (67) into (47) and simple calculation give

$$N = -\sqrt{\frac{6}{h}} \ln \left(\sqrt{\frac{2h\rho_0}{3g + 4h\rho_0}} + \sqrt{\frac{3(g + 2h\rho_0)}{3g + 4h\rho_0}} \right). \quad (71)$$




Figure 1. Left panel. Time evolution of a deep dark soliton in a BEC with repulsive two- and three-body interactions obtained from numerical simulation of the cubic-quintic GPE with $g = 1.0$, $h = 0.5$ and parabolic potential $V(x) = \frac{1}{2}\omega_0^2 x^2$ with $\omega_0 = 0.894$ (the potential is depicted at the top of the figure for scale comparison). Right panel. Dark soliton center coordinate (bottom curve) and soliton density background corresponding to the dynamics depicted in the right panel. The numerical oscillation frequency of the soliton is $\omega = 0.607$, while the time averaged background density, indicated by the dashed line in the figure, is $\langle \rho_0 \rangle = 3.674$. The thin dotted line at the bottom refers to the dynamics of the center of mass of the whole condensate.

Similar calculation of (44) yields

$$m_s = -\sqrt{\frac{6}{h}} \left[\ln \left(\sqrt{\frac{2h\rho_0}{3g+4h\rho_0}} + \sqrt{\frac{3(g+2h\rho_0)}{3g+4h\rho_0}} \right) + \frac{\sqrt{6h\rho_0(g+2h\rho_0)}}{3g+4h\rho_0} \right]. \quad (72)$$

Hence the effective mass of the soliton motion is equal to

$$m_* = 1 + \frac{\sqrt{6h\rho_0(g+2h\rho_0)}}{(3g+4h\rho_0) \ln \left(\sqrt{\frac{2h\rho_0}{3g+4h\rho_0}} + \sqrt{\frac{3(g+2h\rho_0)}{3g+4h\rho_0}} \right)}. \quad (73)$$

For $h\rho_0 \ll g$ this formula reproduces

$$m_* = 2, \quad h\rho_0 \ll g, \quad (74)$$

in agreement with (6), and for $h\rho_0 \gg g$ we get

$$m_* = 1 + \frac{\sqrt{3}}{2 \ln \frac{1+\sqrt{3}}{\sqrt{2}}}, \quad h\rho_0 \gg g, \quad (75)$$

in agreement with (9).

To check these results, we have numerically integrated the GP equation for this case using as initial condition the exact dark soliton solution of the cubic-quintic NLS




Figure 2. Dependence of the ratio $\omega_0/\omega = \sqrt{m^*}$ on the BEC background density ρ_0 for repulsive interactions, $g = 1.0$, $h = 0.5$, (left panel) and for opposite sign interactions $g = 1.0$, $h = -0.5$ (right panel). Continuous curves refer to the analytical results in Eqs. (73), (82), while the dots are obtained from direct numerical solution of the generalized NLS equation (4) with $f(\rho)$ given by Eq. (66).

in Eq. (1). The parabolic potential was introduced adiabatically by rising with time the frequency ω_0 from zero (pure cubic-quintic solution) up to a desired value. Effects of adiabatic changes of parameters on BEC bright and dark solitons were considered in Ref. [22] and the effectiveness of the method was demonstrated. In Fig. 1 we depict the dynamics of a repulsive condensate with its dark soliton inside in presence of a parabolic trap raised adiabatically as described above. In the right panel of this figure we show the dynamics of the center position $X(t)$ of the dark soliton and of the center of mass of the whole condensate $X_{cm} = \int x|\psi|^2 dx$ together with the time evolution of the background density at the soliton position. We see that both the position of the soliton and of the BEC center of mass are very regular and relatively small amplitude excitations develop on the background during the evolution. Note that the center of mass of the condensate oscillates with the trap frequency in agreement with Ehrenfest theorem, while the oscillation frequency and the effective mass of the dark soliton are in very good agreement with our analytical predictions $\omega = 0.599$, $m^* = 2.227$ (for the comparison with numerics we take as ρ_0 the time average of the soliton background measured during the evolution). A more extended comparison between the theory and numerical results is presented in the left panel of Fig. 2 in which we see that in general there is a good qualitative agreement which becomes also quantitatively good at higher values of soliton backgrounds. The deviation observed at lower values of ρ_0 can be ascribed, probably, to a large soliton width comparable with the size of the whole condensate. In this situation the conditions of validity of the theory are not satisfied well enough.

3.2. Attractive quintic interaction ($h < 0$)

For $h < 0$ the integrand function in Eq. (47) is real for

$$\rho_0 < \frac{g}{2|h|}, \quad (76)$$

and this inequality coincides with the condition of modulation stability of BEC, namely the condition that the sound velocity $c_s = \sqrt{\rho_0 f'(\rho_0)}$ is real. It is worth noticing that this condition is more restrictive than the condition $\rho_0 < g/|h|$ that the nonlinearity function is positive, $f(\rho) = g\rho - |h|\rho^2 > 0$. Obviously, we can discuss the soliton oscillations for BECs satisfying the condition (76) only.

In this case $Q(\rho)$ has the double zero ρ_0 and the simple zeroes $\rho_+ > \rho_0$ and $0 < \rho_- < \rho_0$ where

$$\rho_{\pm} = \pm \sqrt{\left(\frac{3g}{4|h|} - \rho_0\right)^2 - \frac{3v^2}{2|h|}} + \left(\frac{3g}{4|h|} - \rho_0\right). \quad (77)$$

Hence there exist both dark and bright soliton solutions. We shall consider here motion only of the dark soliton with the density profile

$$\rho(\xi) = \rho_0 - \frac{(\rho_0 - \rho_-)(\rho_+ - \rho_0)}{(\rho_+ - \rho_-) \cosh^2(\kappa\xi) - (\rho_0 - \rho_-)}, \quad (78)$$

where

$$\kappa = \sqrt{2|h|(\rho_0 - \rho_-)(\rho_+ - \rho_0)/3}. \quad (79)$$

Now calculation of the number of particles N and the soliton mass m_s gives

$$N = -\sqrt{\frac{6}{|h|}} \arcsin \sqrt{\frac{2|h|\rho_0}{3g - 4|h|\rho_0}} \quad (80)$$

and

$$m_s = -\sqrt{\frac{6}{|h|}} \left[\arcsin \sqrt{\frac{2|h|\rho_0}{3g - 4|h|\rho_0}} + \frac{2\sqrt{6|h|\rho_0(g - 2|h|\rho_0)}}{3g - 4|h|\rho_0} \right]. \quad (81)$$

Hence the effective mass of the soliton motion is equal to

$$m_* = \frac{m_s}{N} = 1 + \frac{\sqrt{6|h|\rho_0(g - 2|h|\rho_0)}}{(3g - 4|h|\rho_0) \arcsin \sqrt{\frac{2|h|\rho_0}{3g - 4|h|\rho_0}}}. \quad (82)$$

The condition (76) means that in these expressions the argument of the arcsin function is less than unity. (Of course, expressions (80)–(82) can be obtained from (71)–(73) by means of their analytical continuation to negative values of h .)

For $|h|\rho_0 \ll g$ we again reproduce (74), as it should be. For $g - 2|h|\rho_0 \ll g$ we find the limiting behavior

$$m_* = 1 + \frac{2}{\pi\sqrt{g}} \sqrt{g - 2|h|\rho_0}, \quad g - 2|h|\rho_0 \ll g. \quad (83)$$

Thus, m_* as a function of ρ_0 has a branching point singularity at $\rho_{0c} = g/(2|h|)$ where the system becomes modulationally unstable.

The plot of the ratio of the trap frequency to the soliton's oscillations frequency is illustrated in Fig. 2 (see right panel) and it demonstrates a very good agreement with the results of the numerical simulations.

4. Conclusion

In this paper we have considered the dynamics of a dark soliton inside a BEC described by the quasi-one dimensional GP equation with a parabolic trapping potential. In particular, we have developed the theory which permits one to study the motion of a dark soliton through the stationary background state of BEC and to derive the formulae for the frequency of the oscillation of deep solitons which are very effective for arbitrary form of the nonlinearity function $f(\rho)$. When the form of $f(\rho)$ is appropriately specified, the formulae for the frequency of oscillations reproduce the results of all previously studied special cases. We applied our theory to the study of the practically interesting case of cubic-quintic nonlinearity function $f(\rho) = g\rho + h\rho^2$ and we have demonstrated, as a new effect, the dependence of the frequency of the oscillations on the background density of BEC. In principle this effect can be used for the experimental study of higher nonlinearity on BEC's dynamics.

Acknowledgments

AMK thanks the Department of Physics “E.R. Caianiello” of the University of Salerno, where part of this work was done, for the hospitality received and for financial support. He was also partially supported by RFFI (grant 09-02-00499). MS acknowledges partial support from a MIUR-PRIN initiative.

References

- [1] Pitaevskii L P and Stringari S, *Bose-Einstein Condensation*, (Cambridge University Press, Cambridge, 2003).
- [2] Dalfovo F, Giorgini S, Pitaevskii L P and Stringari S 1999 *Rev. Mod. Phys.* **71** 463
- [3] Matthews M R, Anderson B P, Haljan P C, Hall D S, Wieman C E, Cornell E A 1999 *Phys. Rev. Lett.* **83** 2498
- [4] Strecker K E, Partridge G B, Truscott A G, Hulet R G 2002 *Nature* **417** 150
- [5] Burger S, Bongs K, Dettmer S, Ertmer W, Sengstock K, Sanpera A, Shlyapnikov G V, Lewenstein M 1999 *Phys. Rev. Lett.* **83** 5198
- [6] Denschlag J, Simsarian J E, Feder D L, Clark C W, Collins L A, Cubizolles J, Deng L, Hagley E W, Helmerson K, Reinhart W P, Rolston S L, Schneider B I, Phillips W D 2000 *Science* **287** 97
- [7] Hoefer M A, Ablowitz M J, Coddington I, Cornell E A, Engels P and Schweikhard V 2006 *Phys. Rev. A* **74** 023623
- [8] Kamchatnov A M, Gammal A, and Kraenkel R A 2004 *Phys. Rev. A* **69** 063605
- [9] Pérez-García V M, Michinel H, and Herrero H 1998 *Phys. Rev. A* **57** 3837
- [10] Salasnich L, Parola A and Reatto L, 2002 *Phys. Rev. A* **65** 043614
- [11] Kamchatnov A M and Shchesnovich V S 2004 *Phys. Rev. A* **70** 023604
- [12] Tsuzuki T 1971 *J. Low Temp. Phys.* **4** 441
- [13] Busch Th and Anglin J R 2000 *Phys. Rev. Lett.* **84** 2298

- [14] Frantzeskakis D J, Proukakis N P and Kevrekidis P G 2004 *Phys. Rev. A* **70** 015601
- [15] Konotop V V and Pitaevskii L P 2004 *Phys. Rev. Lett.* **93** 240403
- [16] Brazhnyi V A, Konotop V V and Pitaevskii L P 2006 *Phys. Rev. A* **73** 053601
- [17] Pelinovsky D E, Frantzeskakis D J and Kevrekidis P G 2005 *Phys. Rev. E* **72** 016615
- [18] Frantzeskakis D J, Kevrekidis P G, Proukakis N P 2007 *Phys. Lett. A* **364** 129
- [19] Pelinovsky D E and Kevrekidis P G 2008 *AMS Contemporary Math* **473** 159
- [20] Becker Ch, Stellmer S, Soltan-Panahi P, Dörscher S, Baumert M, Richter E-M, Kronjäger J, Bongs K and Sengstock K 2008 *Nature Phys.* **4** 496
- [21] Kivshar Yu S and Luther-Davies B 1998 *Phys. Rep.* **298** 81
- [22] Abdullaev F Kh and Salerno M 2003 *J. Phys. B: At. Mol. Phys.* **36** 2851